

Sophia Breslin

Thomas Desvignes & John H. Postlethwait



What controls sex
development in fish?

Generally, male and female fish are indistinguishable in appearance.

Antarctic marbled notothenns

Male



Female



In other cases, males and females may look slightly different, making them **sexually dimorphic**.

Antarctic painted notothenns



Striped anal fin



Black anal fin

And sometimes, males and females of the same species are so distinct that previously they were thought to be different species!



Bluehead wrasses



Beyond physical appearance, males and females differ in other ways, especially having different **gonads**—organs that make **gametes**: eggs and sperm.

Females have **ovaries** that produce large **eggs**.

Males have **testes** that produce microscopic **spermatozoa**!






During reproduction, a sperm fuses with an egg to form a **zygote**,

which develops into an **embryo**,

that hatches into a **juvenile**,


that becomes an adult as it develops mature ovaries or testes.



What factors cause gonads to become ovaries or testes?




For most vertebrates, the sex of an animal is determined by its genes, we call that Genetic Sex Determination or **GSD**. A major Sex Determining Gene (**SDG**) on a sex chromosome regulates ovary or testis development.




The **SDG** in mammals is **SRY**, located on the Y chromosome. Individuals with an X and a Y chromosome usually develop testes.




Individuals with two X chromosomes usually develop ovaries.



Birds have the opposite system: their **SDG** is **DMRT1** located on the Z chromosome. Males have two Z chromosomes and two **DMRT1** genes.



Females have one Z and one W chromosome, and so have only one **DMRT1** gene.




Like mammals, many fish have an XX-XY genetic sex determination system...

XX
♀



XY
♂



Humphead cichlids

Goldfin herring cichlids

ZW
♀




ZZ
♂

Some other fish have a ZZ-ZW chromosome system like birds.

Sailfin icefish

$X_1X_1X_2X_2$
♀



X_1X_2Y
♂

Some fish have other sex chromosome variations. For example, males of several Antarctic fish species have a Y chromosome formed by the fusion of a sex-chromosome with a non sex-chromosome.

Strangely, fish have a variety of different SDGs rather than a single gene, like in all mammals or in all birds.

Striped catfish



amhr2

Northern pike




Herring



amh

Rainbow trout



irf9

bmpr1b

Mexican blind cavefish



gdf6

Japanese ricefish



dmy

Greater amberjack



hsd17b1

Luzon ricefish



gsdf

Senegalese sole



fshr

Indian blue ricefish



sox3

But sometimes, even closely related fish have different SDGs, like these three ricefishes.

Some fish even change sex as adults, we call them sequential hermaphrodites.

Protogyny refers to fish that first develop as females and later transition to males. "Proto-" means "first" and "-gyny" refers to "female".




Green birdmouth wrasses



Conversely, **protandry** refers to species that first become males and later transition to females. Here, "-andry" refers to "male", thus "male first".

Clownfish



There are more atypical ways the sex of a fish is determined—or not...


In some species, individuals can alternate between being males or females: a bidirectional sex change.

Bluebanded gobies



In a few species, an individual fish can even have both ovaries and testes and produce eggs and sperm at the same time! They are simultaneous hermaphrodites.


Remotely Operated Vehicles (ROVs) can explore the ocean's depths and observe such rare fish!



Abyssal tripodfish

In some cases, genes don't determine the sex of a fish, instead their environment determines their sex, we call that Environmental Sex Determination or **ESD**.

For example, sometimes temperature can determine the sex of a fish.




Umbrella cichlids

And finally, genes and environment can interact: in some species, the effect of the environment can overrule the **SDG**; this system is referred to as **GSD + ESD**.



In the Northern Gulf of Maine, sex determination in Atlantic silverside is mostly genetic.



But further south, silversides are highly sensitive to temperature. Cool spring water favors development of females, giving them a long time to grow until breeding, while warm summer water favors development of males, who can produce millions of microscopic sperm even with small bodies.

And those are not all the ways by which sex is determined in fish. Almost anything you can imagine exists!

Why are fish sex determination mechanisms so diverse while all birds and all mammals seem to just have one mechanism?



Have changes in sex determination systems contributed to the diversification of fish species? We have much to learn.

Translation credits

Danish: Henrik Lauridsen
French: Thomas Desvignes & Guillaume Lecointre
German: Michael Matschiner & Angelika Schartl
Italian: Luca Schiavon & Chiara Papetti
Norwegian: Benedicte Garmann-Aarhus
Portuguese: Isabela Lagana Ohara & Brian Sidlauskas
Spanish: Manuel Novillo & Alejandro Valdivieso

Font is OpenDislexic-Alta,
a typeface designed for some common symptoms of dyslexia.

This comic was created as part of the
University of Oregon Science and Comics Initiative.

This material is based upon work supported by the Office of Polar Programs
at the National Science Foundation under NSF grant number OPP-2232891.
Any opinions, findings, and conclusions or recommendations expressed
in this material are those of the authors and
do not necessarily reflect the views of the National Science Foundation.

This work was supported by the National Institute of General Medical Sciences (NIGMS)
of the National Institutes of Health under award number R35GM139635.
The content is solely the responsibility of the authors and does not necessarily
represent the official views of the National Institutes of Health.



