

Representations and Sunspot Stability

George W. Evans
University of Oregon

Bruce McGough
Oregon State University

January, 2007

Abstract

By endowing his agents with simple forecasting models, or *representations*, Woodford (1990) found that finite state Markov sunspot equilibria may be stable under learning. We show that common factor representations generalize to all sunspot equilibria the representations used by Woodford (1990). We find that if finite state Markov sunspots are stable under learning then *all* sunspots are stable under learning, provided common factor representations are used.

JEL classification: E32, D83, D84.

Keywords: Indeterminacy; Sunspot Equilibria; E-stability.

1 A brief history of stable sunspots

Sunspot equilibria provide avenues through which agents' expectations can drive fluctuations in real economic activity. Interest in these equilibria developed through the work of Shell (1977), Azariadis (1981), Cass and Shell (1983) and Guesnerie (1986), but remained couched primarily in the theoretical literature until Benhabib and Farmer (1994) and Farmer and Guo (1994) demonstrated the existence of sunspot equilibria in RBC-type models modified to incorporate externalities or monopolistic competition: see Farmer (1999) for a detailed development. These authors, and many other since, have used calibrated DSGE models to argue that fluctuations in agents' expectations explain at least part of the business cycle. These arguments have been extended to New Keynesian monetary models: Clarida, Gali, and Gertler (2000) and Lubik and Schorfheide (2004) suggest that passive monetary policy in the seventies produced an economic environment conducive to sunspot equilibria and the associated high volatility.

The simple existence of sunspot equilibria in a model does not imply their relevance: it may not be possible for agents to coordinate their behavior appropriately. A benchmark coordination device in macroeconomics is stability under learning: for details see Evans and Honkapohja (2001).

In an OLG model, Woodford (1990) found that finite state Markov sunspots may be stable under learning, thus lending credence to the relevance of sunspots for applied models, and in part inspiring the work of Farmer and others. However, Evans and Honkapohja (2001) found that the sunspots studied by Farmer and Guo were not stable under learning. This finding was further supported by Evans and McGough (2005a) and Duffy and Xiao (2005) who searched for stable sunspots in a host of RBC-type models and found none.

The stability of equilibria may depend on the values of the model's parameters; this has been known since Bray and Savin (1986). However, that Woodford's sunspots are stable and Farmer's sunspots are not cannot be explained so easily; indeed, as is well known and shown below, in Woodford's model, Farmer's sunspots are *never* stable. Perhaps then the explanation lies in the stochastic properties of the equilibria in question. Woodford considers equilibria that follow a finite state Markov chain, and Farmer's equilibria are autoregressive with conditional noise captured by a martingale difference sequence with continuous support. However, as we will see below, the stochastic nature of an equilibrium has no impact on its stability.

The stability of sunspot equilibria turns on the way they are viewed by private agents. More formally, a given equilibrium may often be identified with a particular recursion: for example, Farmer considers equilibria in AR(1) form. We called these recursions *representations*. When investigating stability under learning, a representation specifies a natural functional form for the forecasting model agents estimate and use to form their expectations. In Evans and McGough (2005b), we showed for a model with sunspots that a given equilibrium may have several representations consistent with it, and further, that stability under learning is representation dependent. In particular, we found that sunspot equilibria previously thought to be unstable under learning become stable if agents use a forecasting model consistent with a *common factor representation*.

This notion of representation allows us to fully investigate why, in Woodford's model, Farmer's sunspots are never stable and Woodford's sunspots sometimes are. In this paper we show that the representation used in Woodford's analysis is, in fact, a special case of a common factor representation; indeed, common factor representations generalize to all sunspot equilibria the learning mechanism used by Woodford for finite state Markov processes. We conclude that whenever these finite state Markov sunspots are stable under learning, all sunspot equilibria will be, provided a common factor representation is used.

2 Woodford's model: a linearization

The non-stochastic linearized version of Woodford's model is sufficient for our purposes, and is given by

$$y_t = \beta E_t y_{t+1}, \tag{1}$$

where $y_t \in \mathbb{R}$. A rational expectations equilibrium (REE) is any bounded stochastic process y_t satisfying (1). We consider only doubly-infinite processes.

Let y_t be an REE, and set $\varepsilon_t = y_t - E_{t-1}y_t$. Then y_t satisfies the recursion

$$y_t = \beta^{-1}y_{t-1} + \varepsilon_t. \quad (2)$$

We call this recursion the *general form representation* of the equilibrium y_t . Because y_t is bounded we know that either $\varepsilon_t = 0$ (so that $y_{t+k} = 0$ for $k \in \mathbb{Z}$) or $|\beta| > 1$; in the latter case, ε_t can be any martingale difference sequence (m.ds). For the remainder of the paper we assume $|\beta| > 1$. Note that y_t is an REE of (1) if and only if there exists an m.ds ε_t so that y_t satisfies (2). The m.ds ε_t captures variation in y_t due to fluctuations in agents' expectations; it is often called a sunspot and the associated REE y_t is often called a sunspot equilibrium.

To analyze stability under learning, private agents are given a forecasting model whose functional form is consistent with a representation of the REE; this forecasting model is sometimes called a *perceived law of motion* or PLM. For example, to study the stability of an REE y_t , agents might be given the PLM $y_t = \theta'X_t$, where $X_t = (1, y_{t-1}, \varepsilon_t)'$ is the vector of regressors.¹ Using this PLM, agents form expectations which are then imposed into (1) to determine the actual behavior of y_t . The equation identifying this behavior is called the *actual law of motion* or ALM, and under appropriate conditions, it takes the same functional form as the PLM: $y_t = T(\theta)'X_t$. Note that the map $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ takes the perceived coefficients to the actual coefficients, and an REE may be identified with a fixed point θ^* of this map.

Using the T-map, we may write down the ordinary differential equation $\dot{\theta} = T(\theta) - \theta$; a rest point θ^* of this ode corresponds to an REE of (1). We say that the REE is *E-stable* if it is a (locally asymptotically) stable fixed point of the ode. Note that the stability of the fixed point obtains if the eigenvalues of the T-map's derivative, DT , have real part less than one.² We focus on E-stability of equilibria because of the *E-stability Principle*, which states that E-stable REE are locally learnable under least squares or related learning algorithms. The E-stability Principle is known to apply to many models, including (1). For details on the E-stability Principle and the deep connection between E-stability and stochastic convergence of statistical learning algorithms, see Evans and Honkapohja (2001).

3 Results

Let y_t be an REE of (1), and consider its stability by providing agents with a PLM consistent with the representation (2). Straightforward computation shows that DT has 2 as one of its eigenvalues. We conclude with the well-known result that y_t is not stable under learning, and since y_t was arbitrary, no sunspot equilibria of (1) are learnable – at least if agents use a PLM consistent with general form representations.

¹This PLM includes a constant to capture the fact that the linearized model has been recentered at its deterministic steady state – a value not assumed known by agents. Also, normally the PLM would include a perceived error; including this error would be distracting and would not alter our results.

²In case the stability of sunspot equilibria in linear models is being analyzed, the appropriate E-stability condition is that the eigenvalues of DT be less than *or equal* to unity: for details see Evans and Honkapohja (2001).

Equilibria generated by coordination on an arbitrary mds were the type studied by Farmer and others in applied models; however, Woodford had a different type of sunspot in mind. Take as primitive a 2-state Markov process $s_t \in \{0, 1\}$, with transition matrix π . For any $y \in \mathbb{R}^2$, we may construct the associated Markov process

$$y_t = y_i \Leftrightarrow s_t = i - 1, \text{ for } i = 1, 2. \quad (3)$$

Evans and Honkapohja (2003) noted that y_t is an REE of (1) if and only if the following two conditions hold:

$$\pi_{11} + \pi_{22} = 1 + \beta^{-1} \quad (4)$$

$$\bar{y}_2(1 - \pi_{11}) = -\bar{y}_1(1 - \pi_{22}). \quad (5)$$

In case y_t is an REE, we call (3) its *natural representation*. These y_t are the 2-state Markov sunspots for the linear model (1) analogous to those studied in Woodford (1990).

To study the stability under learning of these types of equilibria we follow Evans and Honkapohja (2003): we specify a PLM consistent with (3); that is, we assume agents observe s_t and believe the equilibrium of the model is a 2-state Markov process, but we assume they do not know the values of y_t in these two states.³ Instead, we provide agents with perceived values for the states, and thus we may identify their perceptions with points y in \mathbb{R}^2 . For given perceptions y , the PLM is formally given by (3). The associated T-map is $T_N(y) = \beta\pi y$, that is, if agents believe that the economy's states are y then the states will actually be $\beta\pi y$; also, when perceptions and truth coincide, that is, when $T_N(y) = y$, an REE is identified. Since the eigenvalues of DT evaluated at a fixed point are unity and β , we obtain Evans and Honkapohja's conclusion that y_t is stable under learning provided $\beta < -1$. This is the result for the linear model analogous to the celebrated stability result of Woodford (1990).

Above we noted that if y_t is an REE then there is an mds ε_t so that y_t satisfies (2); thus finite state Markov sunspot equilibria must also have general form representations. Also, even in this case, if agents use a forecasting model consistent with (2) then the equilibrium is unstable. We conclude, as did Evans and Honkapohja, that stability under learning of finite state Markov sunspots is representation dependent.

We now turn to the main question of this paper: "What's so special about finite state Markov sunspots?" The answer, of course, is "Nothing." Let y_t be an REE and ε_t the associated mds. Write $\eta_t = (1 - \beta^{-1}L)^{-1}\varepsilon_t$. The *common factor representation* of y_t is $y_t = \eta_t$; for an extended discussion of common factor representations and their relation to minimal state variable solutions, see Evans and McGough (2005b).⁴ We think of η_t as a serially correlated extrinsic noise process on which agents coordinate to form expectations, and thus we call it a *common factor sunspot*.

To study the stability under learning of y_t , we provide agents with the following PLM: $y_t = a + b\eta_t$. Because $E_t y_{t+1} = a + bE_t \eta_{t+1}$ and because $E_t \eta_{t+1} = \beta^{-1}\eta_t$, we find that

³Incorporating an error into the PLM, as discussed above, makes realistic the assumption that agents perceive the economy is a 2-state process even though the observations do not support this perception.

⁴The name "common factor representation" comes from their construction, which may be thought of as obtained by dividing out the common factor $(1 - \beta^{-1}L)$. This construction is less trivial in higher dimensions and with the incorporation of lags into the reduced form model.

$T_{CF}(a, b) = (\beta a, b)$; thus the eigenvalues of DT are one and β . We conclude that if agents use common factor representations to form their forecasting models then y_t is stable under learning provided $\beta < -1$. Notice we obtain precisely the same condition as Woodford did except that, since y_t was arbitrary, our result applies to *all* sunspot equilibria.

While this result generalizes Woodford's stability result to all sunspots, we have yet to establish the connection between common factor representations and natural representations. Recall our primitive assumption that agents believe the economy follows a 2-state Markov process with transition matrix π , where π satisfies (4). Let Γ be the set of all 2-state Markov processes with transition matrix π and notice that Γ may be identified with \mathbb{R}^2 . We think of Γ as the set of all possible agent beliefs.

We may also think of Γ as the set of all PLMs consistent with natural representations: simply recall that s_t is the primitive sunspot with transition matrix π , let $y \in \Gamma$ and see equation (3). While the identification of Γ with this set is, in a sense, trivial, distinguishing the sets will aid clarity; therefore, let Γ_N be the set of all PLMs consistent with natural representations (as captured by \mathbb{R}^2 together with equation (3)) and let $S_N : \Gamma \rightarrow \Gamma_N$ be the identity map on \mathbb{R}^2 . The map S_N , then, takes an agent's beliefs to the associated PLM.

Finally, we must characterize the set of all PLMs consistent with common factor representations. To this end, let y_t be a 2-state Markov sunspot equilibrium with transition matrix π and states $\bar{y} \in \mathbb{R}^2$ satisfying (5), let ε_t be the mds such that y_t solves (2), and let η_t be the common factor sunspot generated by ε_t . Notice that η_t is a 2-state Markov process. Now let Γ_{CF} be the set of PLMs consistent with the common factor representation of y_t , and notice that a PLM is uniquely determined by its coefficients: $a + b\eta_t = c + d\eta_t \Leftrightarrow a = c$ and $b = d$. Recalling that Γ is just \mathbb{R}^2 , we may define $S_{CF} : \Gamma \rightarrow \Gamma_{CF}$ by

$$S_{CF}(y) = \left(\frac{y_1(\bar{y}_2 - \bar{y}_1) - \bar{y}_1(y_2 - y_1)}{\bar{y}_2 - \bar{y}_1}, \frac{y_2 - y_1}{\bar{y}_2 - \bar{y}_1} \right)'.$$

Straightforward computation shows that S_{CF} is a bijection. Finally, define $S : \Gamma_N \rightarrow \Gamma_{CF}$ by $S = S_{CF} \circ S_N^{-1}$ and note that S is bijective by construction.

Proposition 1 *The following diagram commutes:*

$$\begin{array}{ccccc}
 & & \Gamma_N & \xrightarrow{T_N} & \Gamma_N \\
 & \nearrow S_N & \downarrow S & & \downarrow S \\
 \Gamma & \xrightarrow{S_{CF}} & \Gamma_{CF} & \xrightarrow{T_{CF}} & \Gamma_{CF} & \xrightarrow{S_{CF}^{-1}} & \Gamma \\
 & & & & & \nwarrow S_N^{-1} &
 \end{array}$$

A commutative diagram is an efficient way to make statements about equivalence of functions. To understand the diagram's meaning, start with a point in any set (the "initial" set)

and pick any other set (the “final” set) that can be reached from the initial set by following a path of arrows. Under the composition of the functions corresponding to the arrows in your path, the point you chose in the initial set is mapped to a point in the final set. Because the diagram commutes, the path of arrows you chose to reach the final set is irrelevant. An important example is

$$S_N^{-1} \circ T_N \circ S_N = S_{CF}^{-1} \circ T_{CF} \circ S_{CF}. \quad (6)$$

The proof of the proposition is algebra using the restrictions (4) and (5), and is available from the authors upon request.

The commutivity of the above diagram allows us to make several precise statements about the relationship between common factor representations and Woodford’s natural representations.

Corollary 2 *PLMs consistent with common factor representations and PLMs consistent with natural representations identify the same set of agent beliefs.*

Corollary 3 *Viewed as acting on agents’ beliefs, the maps T_N and T_{CF} coincide.*

Corollary 2 acknowledges that S is bijective and Corollary 3 is an interpretation of equation (6). Taken together, these corollaries indicate the sense in which, when restricted to finite state Markov sunspot equilibria, common factor representations may be identified with Woodford’s natural representations; importantly, however, common factor representations may be used to analyze the stability of any sunspot equilibrium. In this sense, we may view common factor representations as a generalization of Woodford’s natural representations to all sunspot equilibria.

Finally, by applying the chain rule and the inverse function theorem to equation (6) we find that the eigenvalues of DT_N and DT_{CF} coincide. Thus

Corollary 4 *Common factor representations and natural representations have the same stability properties.*

Furthermore, because the eigenvalues of the associated T-maps are independent of the cardinality of the sunspot’s support, we may conclude that whenever finite state Markov sunspots are stable under learning, all sunspot equilibria are stable under learning, provided common factor representations are used for the stability analysis.⁵

4 Conclusion

We have shown that common factor representations generalize Woodford’s natural representations and his stability results to all sunspot equilibria. This indicates that Woodford’s

⁵While this conclusion only applies to the simple model considered in this paper, we conjecture that it holds more generally.

stability results may extend to many other models, and to assess this possibility, common factor representations should be analyzed.

References

- AZARIADIS, C. (1981): “Self-Fulfilling Prophecies,” *Journal of Economic Theory*, 25, 380–396.
- BENHABIB, J., AND R. A. FARMER (1994): “Indeterminacy and Increasing Returns,” *Journal of Economic Theory*, 63, 19–41.
- BRAY, M., AND N. SAVIN (1986): “Rational Expectations Equilibria, Learning, and Model Specification,” *Econometrica*, 54, 1129–1160.
- CASS, D., AND K. SHELL (1983): “Do Sunspots Matter?,” *Journal of Political Economy*, 91, 193–227.
- CLARIDA, R., J. GALI, AND M. GERTLER (2000): “Monetary Policy Rules and Macroeconomic Stability: Evidence and Some Theory,” *Quarterly Journal of Economics*, 115, 147–180.
- DUFFY, J., AND W. XIAO (2005): “Instability of Sunspot Equilibria in Real Business Cycle Models under Adaptive Learning,” *Journal of Monetary Policy*, Forthcoming.
- EVANS, G. W., AND S. HONKAPOHJA (2001): *Learning and Expectations in Macroeconomics*. Princeton University Press, Princeton, New Jersey.
- (2003): “Expectational Stability of Stationary Sunspot Equilibria in a Forward-looking Model,” *Journal of Economic Dynamics and Control*, 28, 171–181.
- EVANS, G. W., AND B. MCGOUGH (2005a): “Indeterminacy and the Stability Puzzle in Non-Convex Economies,” *Contributions to Macroeconomics, BE-Press*, 5.
- (2005b): “Stable Sunspot Solutions in Models with Predetermined Variables,” *Journal of Economic Dynamics and Control*, 29, 601–625.
- FARMER, R. E. (1999): *The Economics of Self-Fulfilling Prophecies, Second edition*. MIT Press, Cambridge, Mass.
- FARMER, R. E., AND J.-T. GUO (1994): “Real Business Cycles and the Animal Spirits Hypothesis,” *Journal of Economic Theory*, 63, 42–72.
- GUESNERIE, R. (1986): “Stationary Sunspot Equilibria in an N-commodity World,” *Journal of Economic Theory*, 40, 103–128.
- LUBIK, T., AND F. SCHORFHEIDE (2004): “Testing for Indeterminacy: an application to U.S. Monetary Policy,” *American Economic Review*, 94, 190–217.

SHELL, K. (1977): “Monnaie et Allocation Intertemporelle,” Working paper, CNRS Séminaire de E.Malinvaud, Paris.

WOODFORD, M. (1990): “Learning to Believe in Sunspots,” *Econometrica*, 58, 277–307.